

On the Squeezed Vacuum States Corresponding to the General Two-Mode Bogolubov Transformation

Oleg Mokhov[†]

*Department of Mathematics
Bilkent University
06533 Bilkent, Ankara, Turkey*

Abstract

In this paper the explicit form of the operator of transformation of the vacuum states for the general two-mode Bogolubov transformation is found

[†] On leave of absence from *VNIIFTRI, Mendeleevo, Moscow Region 141570, Russia*

1 Introduction

A number of problems of quantum optics is connected with investigation of model Hamiltonians given by Hermitian quadratic form of Bose operators:

$$H = \frac{1}{2} \sum_{k,l} [A_{kl} a_k^\dagger a_l^\dagger + A_{kl}^* a_k a_l + 2B_{kl} a_k^\dagger a_l], \quad (1)$$

$$A_{kl} = A_{lk}, \quad B_{kl} = B_{lk}^*,$$

$$[a_k, a_l^\dagger] = \delta_{kl}, \quad [a_k, a_l] = [a_k^\dagger, a_l^\dagger]$$

For diagonalization of such quadratic Hamiltonians the canonical Bogolubov transformations [1] are used:

$$b_k = \sum_m (u_{km} a_m + v_{km} a_m^\dagger), \quad (2)$$

$$b_k^\dagger = \sum_m (v_{km}^* a_m + u_{km}^* a_m^\dagger),$$

i.e. the linear transformations of Bose operators, preserving the commutational relations of the algebra of Bose operators:

$$[b_k, b_l^\dagger] = \delta_{kl}, \quad [b_k, b_l] = [b_k^\dagger, b_l^\dagger] \quad (3)$$

In particular, the use of the general canonical Bogolubov transformations is necessary for the study of non-classical optical effects such as squeezing [2,3] (see also the survey [4]). In the case of a single mode, for the Hermitian quadrature operators

$$Q = \frac{(a^\dagger + a)}{2}, \quad P = i \frac{(a^\dagger - a)}{2}, \quad [Q, P] = \frac{i}{2} \quad (4)$$

we have the Heisenberg uncertainty relation for quantum fluctuations:

$$\langle (\Delta Q)^2 \rangle \langle (\Delta P)^2 \rangle \geq \left| \frac{\langle [Q, P] \rangle}{4} \right|^2 \equiv \frac{1}{16} \quad (5)$$

The vacuum and coherent states give

$$\langle (\Delta Q)^2 \rangle = \langle (\Delta P)^2 \rangle = \frac{1}{4},$$

i.e. they realize the minimum-uncertainty states with

$$\langle (\Delta Q)^2 \rangle \langle (\Delta P)^2 \rangle = \frac{1}{16} \quad (6)$$

For squeezed states

$$\langle (\Delta Q)^2 \rangle = \frac{1}{4} \exp(-2r), \quad \langle (\Delta P)^2 \rangle = \frac{1}{4} \exp(2r) \quad (7)$$

and they also realize the minimum-uncertainty states but if $r \neq 0$ then one of the quadrature operators satisfies to the relation $\langle (\Delta X)^2 \rangle < \frac{1}{4}$ ($X = Q$ or $X = P$), i.e. the quantum fluctuations in one of the field quadrature can be below than the usual vacuum level. These squeezed states can be obtained with the use of the general one-mode canonical Bogolubov transformation:

$$b = e^{i\phi} \cosh r \, a + e^{i\psi} \sinh r \, a^\dagger, \quad b^\dagger = e^{-i\phi} \cosh r \, a^\dagger + e^{-i\psi} \sinh r \, a \quad (8)$$

In this connection one needs an explicit form of the operators transforming the Fock states of Bose-fields a_k to the corresponding Fock states of Bose-fields b_k (for Bose-fields a_k and b_k connected by the Bogolubov transformation (2),(3)). First of all, it is necessary to know the operator of transformation for the corresponding vacuum states $a_k|0\rangle^{(a)} = 0$, $b_k|0\rangle^{(b)} = 0$, $1 \leq k \leq N$, where $|0\rangle^{(x)} = \prod_{m=1}^N |0\rangle_m^{(x)}$. For one-mode case (a single Bose-field a) such unitary operator (so called "squeezed" operator) was found by Stoler [2] for the general one-mode canonical Bogolubov transformation (8). For two-mode case analogous "squeezed" operators was constructed only for Bogolubov transformations of the very special form, as a matter of fact, only for the direct combination of one-mode Bogolubov transformations (after some elementary transformations of Bose-fields) [5,6], see also [4]. Meanwhile, for quantum optics the general two-mode case is also very important. Particularly, the model of polariton [7,8] corresponds to the more general case of two-mode quadratic Hamiltonian. In this paper we attempted to study the most general two-mode case. The explicit formulas connecting the general Bogolubov transformation with the corresponding unitary "squeezed" operator are very complicated but we present here the simple explicit form of the operator of transformation for the vacuum states for the most general two-mode Bogolubov transformation.

Author is very grateful to Prof. A.S.Shumovsky for valuable discussions on the problems of quantum optics.

2 General scheme for N-mode case

We shall consider here the N-mode case of the usual operators of creation and annihilation for photons:

$$a_k|n\rangle_k = \sqrt{n}|n-1\rangle_k, \quad a_k^\dagger|n\rangle_k = \sqrt{n+1}|n+1\rangle_k, \quad a_k^\dagger a_k|n\rangle_k = n|n\rangle_k$$

Let be given an arbitrary unitary transformation of Bose operators a :

$$b_k = U a_k U^\dagger, \quad (9)$$

where $U(a, a^\dagger)$ is an unitary operator: $U^\dagger = U^{-1}$, $b_k^\dagger = U a_k^\dagger U^\dagger$. The transformation (9) preserve the commutational relations (3) for the algebra of Bose operators. The vacuum state of Bose-fields b is determined by the following formula:

$$|0 \rangle^{(b)} = U |0 \rangle^{(a)}$$

For arbitrary Fock states we have

$$\begin{aligned} \prod_{k=1}^N |n_k \rangle_k^{(b)} &= \prod_{k=1}^N \frac{(b_k^\dagger)^{n_k}}{\sqrt{(n_k)!}} |0 \rangle_k^{(b)} = \\ U \prod_{k=1}^N \frac{(a_k^\dagger)^{n_k}}{\sqrt{(n_k)!}} |0 \rangle_k^{(a)} &= U \prod_{k=1}^N |n_k \rangle_k^{(a)}, \end{aligned}$$

i.e. the unitary operator U is the operator of transformation for all Fock states after the unitary transformation (9) of Bose operators.

Consider now an arbitrary Bogolubov transformation (2),(3). For the vacuum state of Bose-fields b we have:

$$\begin{aligned} |0 \rangle^{(b)} &= \sum_{i_s=0}^{\infty} \lambda_{i_1 \dots i_N} \prod_{k=1}^N |i_k \rangle_k^{(a)}, \\ 1 \leq s \leq N \end{aligned} \quad (10)$$

The condition

$$b_k |0 \rangle^{(b)} = 0$$

gives recurrence formulas for the coefficients $\lambda_{i_1 \dots i_N}$:

$$\sum_{m=1}^N (u_{km} \lambda_{i_1 \dots (i_m+1) \dots i_N} \sqrt{i_m+1} + v_{km} \lambda_{i_1 \dots (i_m-1) \dots i_N} \sqrt{i_m}) = 0 \quad (11)$$

For $N = 1$ and $N = 2$ these equations for $\lambda_{i_1 \dots i_N}$ are completely solved in this paper for the corresponding general Bogolubov transformations (of course, the case $N = 1$ is very simple). After that if we know the operator $S_0(a^\dagger)$ such that

$$|0 \rangle^{(b)} = S_0(a^\dagger) |0 \rangle^{(a)}$$

we can determine uniquely the corresponding unitary "squeezed" operator U by recurrence procedure from the necessary relations for all photon number Fock states:

$$\prod_{k=1}^N |n_k \rangle_k^{(b)} = U \prod_{k=1}^N |n_k \rangle_k^{(a)}$$

3 One-mode case

First of all consider the simple one-mode case of the general Bogolubov transformation:

$$b = ua + va^\dagger, \quad b^\dagger = v^*a + u^*a^\dagger$$

The condition $[b, b^\dagger] = 1$ gives $uu^* - vv^* = 1$, i.e.

$$u = e^{i\phi} \cosh r, \quad v = e^{i\psi} \sinh r$$

For the vacuum state of Bose-field b we have:

$$|0\rangle^{(b)} = \sum_{i=0}^{\infty} \lambda_i |i\rangle^{(a)}, \quad (12)$$

and from the condition $b|0\rangle^{(b)} = 0$ we receive the simple recurrence formulas for λ_i :

$$\lambda_{i+1}u\sqrt{i+1} + \lambda_{i-1}v\sqrt{i} = 0 \quad (13)$$

It is easy to obtain the general solution of these recurrence equations (13):

$$\lambda_{2k+1} = 0, \quad \lambda_{2k} = (-1)^k \left(\frac{v}{u}\right)^k \frac{\sqrt{(2k-1)!!}}{\sqrt{(2k)!!}} \lambda_0 \quad (14)$$

Thus,

$$\begin{aligned} |0\rangle^{(b)} &= \sum_{k=0}^{\infty} (-1)^k \left(\frac{v}{u}\right)^k \frac{\sqrt{(2k-1)!!}}{\sqrt{(2k)!!}} \lambda_0 |2k\rangle^{(a)} \equiv \\ &\left(\sum_{k=0}^{\infty} (-1)^k \left(\frac{v}{2u}\right)^k \frac{(a^\dagger)^{2k}}{k!} \right) \lambda_0 |0\rangle^{(a)}, \end{aligned} \quad (15)$$

i.e. we receive the following formula for transformation of the vacuum states in the one-mode case:

$$|0\rangle^{(b)} = \lambda_0 e^{-\frac{v}{2u}(a^\dagger)^2} |0\rangle^{(a)} \quad (16)$$

The corresponding unitary operator

$$U = \lambda_0 \sum_{k=0}^{\infty} S_k (a^\dagger)^k a^k, \quad (17)$$

where

$$S_0(a^\dagger) = \exp\left(-\frac{v}{2u}(a^\dagger)^2\right), \quad (18)$$

can be determined uniquely from relations

$$|n \rangle^{(b)} = U |n \rangle^{(a)}$$

which give the recurrence formulas for $S_n(a^\dagger)$:

$$(v^* a + u^* a^\dagger)^n S_0(a^\dagger) |0 \rangle^{(a)} = \sum_{k=0}^{\infty} S_k(a^\dagger) a^k (a^\dagger)^n |0 \rangle^{(a)} \quad (19)$$

For $\phi = 0$ we have the well-known unitary "squeezed" operator of Stoler [2]:

$$U = \exp \left[\frac{1}{2} (r e^{-i\psi} a^2 - r e^{i\psi} (a^\dagger)^2) \right]$$

4 Two-mode case

Consider the general two-mode Bogolubov transformation:

$$\begin{aligned} b_1 &= u_{11} a_1 + u_{12} a_2 + v_{11} a_1^\dagger + v_{12} a_2^\dagger, \\ b_2 &= u_{21} a_1 + u_{22} a_2 + v_{21} a_1^\dagger + v_{22} a_2^\dagger, \\ b_1^\dagger &= v_{11}^* a_1 + v_{12}^* a_2 + u_{11}^* a_1^\dagger + u_{12}^* a_2^\dagger, \\ b_2^\dagger &= v_{21}^* a_1 + v_{22}^* a_2 + u_{21}^* a_1^\dagger + u_{22}^* a_2^\dagger, \end{aligned} \quad (20)$$

where the coefficients (u_{ij}, v_{ij}) must satisfy the relations:

$$\begin{aligned} 1) \quad & u_{11} v_{21} + u_{12} v_{22} - u_{21} v_{11} - u_{22} v_{12} = 0, \\ 2) \quad & u_{11} u_{21}^* + u_{12} u_{22}^* - v_{11} v_{21}^* - v_{12} v_{22}^* = 0, \\ 3) \quad & u_{11} u_{11}^* + u_{12} u_{12}^* - v_{11} v_{11}^* - v_{12} v_{12}^* = 1, \\ 4) \quad & u_{21} u_{21}^* + u_{22} u_{22}^* - v_{21} v_{21}^* - v_{22} v_{22}^* = 1, \end{aligned} \quad (21)$$

in order to preserve the commutational relations (3) of the algebra of Bose operators. For the vacuum state of Bose-fields b we have the following representation:

$$|0 \rangle^{(b)} = \sum_{i,j=0}^{\infty} \lambda_{ij} |i \rangle_1^{(a)} |j \rangle_2^{(a)} \quad (22)$$

and from the conditions $b_k|0 \rangle^{(b)} = 0$ we can obtain the recurrence equations for λ_{ij} :

$$\begin{aligned}\lambda_{(i+1)j}u_{11}\sqrt{i+1} + \lambda_{i(j+1)}u_{12}\sqrt{j+1} + \lambda_{(i-1)j}v_{11}\sqrt{i} + \lambda_{i(j-1)}v_{12}\sqrt{j} &= 0, \\ \lambda_{(i+1)j}u_{21}\sqrt{i+1} + \lambda_{i(j+1)}u_{22}\sqrt{j+1} + \lambda_{(i-1)j}v_{21}\sqrt{i} + \lambda_{i(j-1)}v_{22}\sqrt{j} &= 0\end{aligned}\quad (23)$$

Further we shall consider that

$$\Delta_1 \equiv u_{11}u_{22} - u_{12}u_{21} \neq 0$$

The case $\Delta_1 = 0$ is very simple and we shall not stop here on it. Let us introduce also for convenience the following notations:

$$\Delta_2 \equiv u_{11}v_{21} - u_{21}v_{11},$$

$$\Delta_3 \equiv u_{11}v_{22} - u_{21}v_{12},$$

$$\Delta_4 \equiv u_{22}v_{11} - u_{12}v_{21},$$

$$\Delta_5 \equiv u_{22}v_{12} - u_{12}v_{22}.$$

Now we can transform the system (23) to the form:

$$\begin{aligned}\lambda_{i(j+1)}\Delta_1\sqrt{j+1} &= -\lambda_{(i-1)j}\Delta_2\sqrt{i} - \lambda_{i(j-1)}\Delta_3\sqrt{j}, \\ \lambda_{(i+1)j}\Delta_1\sqrt{i+1} &= -\lambda_{(i-1)j}\Delta_4\sqrt{i} - \lambda_{i(j-1)}\Delta_5\sqrt{j}\end{aligned}\quad (24)$$

The system (23) is compatible if and only if $\Delta_2 = \Delta_5$ but this is exactly the condition that $[b_1, b_2] = 0$ (see the relation 1) (21)), i.e. this relation is always valid for the canonical Bogolubov transformation. Under the unique condition $\Delta_2 = \Delta_5$ we can obtain the general solution of the system (23):

$$\begin{aligned}\lambda_{(2k)(2n+1)} &= \lambda_{(2k+1)(2n)} = 0, \\ \lambda_{(2k)(2n)} &= (-1)^{(n+k)}\sqrt{(2n)!}\sqrt{(2k)!}\times \\ &\sum_{0 \leq s \leq n; s \leq k} \left[\left(\frac{\Delta_2}{\Delta_1} \right)^{2s} \left(\frac{\Delta_3}{2\Delta_1} \right)^{n-s} \left(\frac{\Delta_4}{2\Delta_1} \right)^{k-s} \frac{1}{(n-s)!(k-s)!(2s)!} \right] \lambda_0, \\ \lambda_{(2k+1)(2n+1)} &= (-1)^{(n+k+1)}\sqrt{(2n+1)!}\sqrt{(2k+1)!}\times \\ &\sum_{0 \leq s \leq n; s \leq k} \left[\left(\frac{\Delta_2}{\Delta_1} \right)^{2s+1} \left(\frac{\Delta_3}{2\Delta_1} \right)^{n-s} \left(\frac{\Delta_4}{2\Delta_1} \right)^{k-s} \frac{1}{(n-s)!(k-s)!(2s+1)!} \right] \lambda_0\end{aligned}\quad (25)$$

Thus, for the vacuum state of Bose-fields b we have:

$$|o >^{(b)} = \sum_{k,n} \left[\lambda_{(2k)(2n)} |2k >_1^{(a)} |2n >_2^{(a)} + \lambda_{(2k+1)(2n+1)} |2k+1 >_1^{(a)} |2n+1 >_2^{(a)} \right]$$

or in the operator form:

$$|0 >^{(b)} = \lambda_0 e^{-\frac{\Delta_4}{2\Delta_1}(a_1^\dagger)^2 - \frac{\Delta_2}{\Delta_1} a_1^\dagger a_2^\dagger - \frac{\Delta_3}{2\Delta_1}(a_2^\dagger)^2} |0 >^{(a)} \quad (26)$$

Thus, the operator

$$S_0(a^\dagger) = \exp \left(-\frac{\Delta_4}{2\Delta_1}(a_1^\dagger)^2 - \frac{\Delta_2}{\Delta_1} a_1^\dagger a_2^\dagger - \frac{\Delta_3}{2\Delta_1}(a_2^\dagger)^2 \right) |0 >^{(a)} \quad (27)$$

is the operator of transformation of the vacuum states for the most general two-mode Bogolubov transformations (20),(21). The corresponding unitary "squeezed" operator U can be determined now by recurrence procedure from the conditions

$$|n_1 >_1^{(b)} |n_2 >_2^{(b)} = U |n_1 >_1^{(a)} |n_2 >_2^{(a)}$$

References

- [1] N.N.Bogolubov, J.Phys.USSR 11(1947), 23
- [2] D.Stoler, Phys. Rev. D. 1(1970), 3217; D. 4(1971), 1925
- [3] H.P.Yuen, Phys.Rev. A. 13(1976),2226
- [4] R.Loudon, P.L.Knight, J.Modern Optics 34(1987), 709
- [5] G.J.Milburn, J.Phys. A. 17(1984), 737
- [6] C.M.Caves, B.L.Schumaker, Phys.Rev. D. 31(1985), 3068
- [7] A.S.Davydov, Theory of solid. Nauka, Moscow. 1971
- [8] A.V.Chizhov, R.G.Nazmitdinov, A.S.Shumovsky, Quant. Optics 3(1991), 1